

Get them IN ON TIME

Birth	Age 2 months Not earlier than 6 weeks of age	Age 4 months Interval from previous dose	Age 6 months Interval from previous dose	Age 12 months Interval from previous dose	Age 4-6 YEARS Before Kindergarten	Age 11-12 YEARS Middle School
	RV (Rotavirus)	RV (Rotavirus)	RV (Rotavirus)			
HepB	HepB		HepB at least 4 months after first dose			
	DTaP	DTaP 1-2 months	DTaP 1-2 months	DTaP 6 months after the third dose	DTaP	Tdap
	Hib Not earlier than 6 weeks of age	Hib 1-2 months	Hib 1-2 months	Hib 1-2 months		
	Polio	Polio 1-2 months	Polio 1-2 months		Polio	
	PCV	PCV 1-2 months	PCV 1-2 months	PCV 6-8 weeks and on or after first birthday		
				MMR On or after first birthday	MMR	
				Varicella On or after first birthday	Varicella	
				HepA 2nd dose 6 months later		
						MCV4 2 shot series
						HPV 3 shot series

Every Fall: Flu Vaccine

• All children 6 months to 18 years of age.

Note: Children 6 months through 8 years old need 2 doses, one month apart, in their first year of flu shots.

Any questions please call (260) 424-SHOT or visit: supershot.org

Indiana's "IN ON TIME" Childhood Immunization Schedule is compatible with the recommendation of the Advisory Committee on Immunization Practices (ACIP) of the US Centers for Disease Control and Prevention (CDC), the American Academy of Pediatrics (AAP) and the American Academy of Family Physicians (AAFP). If you have any questions please call the CDC at 800-232-4636 (800-CDC-INFO).

Combined vaccines may decrease the number of shots.